

Experimental results on $(sd)^3$ structures of ^{16}C and ^{17}C

Three-neutron $(sd)^3 \otimes$ core configurations

H.G. Bohlen¹, R. Kalpakchieva², W. von Oertzen¹, T.N. Massey³, A.A. Ogloblin⁴
G. de Angelis⁵, M. Milin⁶, Ch. Schulz¹, Tz. Kokalova¹ and C. Wheldon¹

¹ Hahn-Meitner-Institut, Berlin, Germany

² FLNR, JINR, Dubna, Russia

³ Ohio University, Athens, USA

⁴ IGNP, Kurchatov Institute, Moscow, Russia

⁵ LNL, INFN, Legnaro, Italy

⁶ Rudjer Boskovic Institute, Zagreb, Croatia

Ground state properties and first excited states of ^{16}C and ^{17}C

^{16}C : States of ^{16}C known up to 6.11 MeV excitation energy from the $^{14}\text{C}(t,p)$ reaction [Balamuth77, Fortune77, Sercely78]. Well described by shell model (SM) calculations. States up to 17.4 MeV observed in the $^{13}\text{C}(^{12}\text{C},^9\text{C})$ reaction [Boh03].

Ground state properties were investigated by one-neutron removal reactions using a ^{16}C radioactive beam [many refs., see, e.g., Maddalena01].

Result: 58(6) % $(1d5/2)^2$ and 42(6) % $(2s1/2)^2$ in good agreement with SM calculs.

B(E2) value of the first 2^+ state at 1.77 MeV is extremely small [Imai04, Elekes04].

Ground state properties and first excited states of ^{16}C and ^{17}C

^{16}C : States of ^{16}C known up to 6.11 MeV excitation energy from the $^{14}\text{C}(t,p)$ reaction [Balamuth77, Fortune77, Sercely78]. Well described by shell model (SM) calculations. States up to 17.4 MeV observed in the $^{13}\text{C}(^{12}\text{C},^9\text{C})$ reaction [Boh03].

Ground state properties were investigated by one-neutron removal reactions using a ^{16}C radioactive beam [many refs., see, e.g., Maddalena01].

Result: 58(6) % $(1d5/2)^2$ and 42(6) % $(2s1/2)^2$ in good agreement with SM calculs.

B(E2) value of the first 2^+ state at 1.77 MeV is extremely small [Imai04, Elekes04].

^{17}C : Only two states of ^{17}C observed in the $^{48}\text{Ca}(^{18}\text{O},^{17}\text{C})^{49}\text{Ti}$ reaction [Nolen77, Fifield82], the ground state and an excited state at 0.295(10) MeV (probably $5/2^+$).

Puzzle: the spin-parity of the ^{17}C ground state: ?? $1/2^+$, $3/2^+$, $5/2^+$??

Solution: **→ $3/2^+$!!**

a) β -decay of ^{17}C to exc. states of ^{17}N and γ -decay branching ratios:

$5/2^+$ excluded [Warburton, Millener89].

b) g-factor measurement [Ogawa02]: $1/2^+$ excluded


c) ^{17}C -beam, 1n-removal cross sections confirm $3/2^+$ [Maddal.01, Sauvan04, Datta03]

Recently γ -transitions were reported at 0.21 MeV and 0.33 MeV [Stanoiu04]

Low-lying states of ^{16}C and ^{17}C :

^{16}C : $^{14}\text{C}(\text{gs}) \otimes \nu(\text{sd})^2$


Schematic $(\text{sd})^2$:




Low-lying states of ^{16}C and ^{17}C :


^{16}C : $^{14}\text{C}(\text{gs}) \otimes \nu(\text{sd})^2$ and $^{17}\text{C}, ^{16}\text{C}$: $^{14,13}\text{C}(\text{gs}) \otimes \nu(\text{sd})^3$, respectively.

Schematic $(\text{sd})^2$: $(\text{sd})^3$: direct population of three-neutron configurations



ISL at HMI Berlin










Reaction conditions and selectivity

The $^{12}\text{C}(^{12}\text{C}, ^9\text{C})^{15}\text{C}$ - reaction at 231 MeV

- Breit-Wigner resonances ($S_n = 1.22\text{ MeV}$)
- Three-body distributions $^9\text{C} + n + ^{14}\text{C}^*$
 sequential decay: $^{10}\text{C}^*(22.2\text{MeV}) (3^-, 6.73\text{MeV})$






Reaction conditions and selectivity

The $^{12}\text{C}(^{12}\text{C}, ^9\text{C})^{15}\text{C}$ - reaction at 231 MeV

- Breit-Wigner resonances ($S_n = 1.22\text{ MeV}$)
- Three-body distributions $^9\text{C} + n + ^{14}\text{C}^*$
 sequential decay: $^{10}\text{C}^*(22.2\text{MeV}) (3^-, 6.73\text{MeV})$
- Cross sections are strongly dependent on dynamical matching conditions:
 large negative Q-values \rightarrow extreme mismatch of grazing angular momenta: $L_{gr,i} - L_{gr,f} \geq 9!$
 \Rightarrow stretched configurations are strongly favored
 e.g. $9/2^- : [(1d5/2)_{4+}^2 \times (1p1/2)^{-1}] 9/2^-$ yes ($\uparrow\uparrow$)
 $7/2^- : [(1d5/2)_{4+}^2 \times (1p1/2)^{-1}] 7/2^-$ no ($\uparrow\downarrow$)
 $\Rightarrow \ell=0$ angular momenta are strongly disfavored





Spectroscopy of ^{16}C using the three-neutron transfer ($^{12}\text{C}, ^9\text{C}$)

^{15}C
 (^{12}C -target, two holes in the 1p-shell)
 energy resolution: 0.20 - 0.25 MeV

^{16}C :
 (^{13}C -target, core with 1p1/2-hole built-in)


14 new states observed [Boh03]
 states known from (t,p) [e.g., Fortune78]





Spectroscopy of ^{16}C using the three-neutron transfer ($^{12}\text{C}, ^9\text{C}$)

^{15}C
 (^{12}C -target, two holes in the 1p-shell)
 energy resolution: 0.20 - 0.25 MeV




^{16}C :
 (^{13}C -target, core with 1p1/2-hole built-in)

14 new states observed [Boh03]
 known states

stretched configurations are **strong**:
 $4^+ [(1d5/2)^2]$
 $5^- [(1d5/2)^3, 9/2^+ \times (1/2^-)^{-1}]$

$5^-, 4^-$ doublet: $9/2^+ \uparrow\uparrow 1/2^-, 9/2^+ \uparrow\downarrow 1/2^-$
 $3^-, 2^-$ doublet: $5/2^+ \uparrow\uparrow 1/2^-, 5/2^+ \uparrow\downarrow 1/2^-$



^{14}C -target, background subtracted


$S_n = 0.73 \text{ MeV}$, $Q_0 = -46.93 \text{ MeV}$

13 new states identified above S_n

strongest state: stretched config. $(1d5/2)^3 \rightarrow 9/2^+$

Starting with: $^{14}\text{C}_{\text{gs}}$ target (closed neutron 1p-shell)
 direct 3n-transfer to the open sd-shell

\Rightarrow population of $(sd)^3$ structures expected ($3p-0h$)



^{14}C -target, background subtracted


$$S_n = 0.73 \text{ MeV}, \quad Q_0 = -46.93 \text{ MeV}$$

13 new states identified above S_n

strongest state: stretched config. $(1d5/2)^3 \rightarrow 9/2^+$


Starting with: $^{14}\text{C}_{\text{gs}}$ target (closed neutron 1p-shell)
 direct 3n-transfer to the open sd-shell

\Rightarrow population of $(sd)^3$ structures expected ($3p-0h$)



Comparison of ^{17}C - ^{16}C spectra

Spectra aligned for the states with $(1d5/2)^3, 9/2^+$ configurations, in ^{16}C at 8.92 MeV and in ^{17}C at 3.10 MeV. (offset 5.82 MeV, same scale factor)



^{14}C -target, background subtracted


$S_n = 0.73 \text{ MeV}$, $Q_0 = -46.93 \text{ MeV}$

13 new states identified above S_n

strongest state: stretched config. $(1d5/2)^3 \rightarrow 9/2^+$

Starting with: $^{14}\text{C}_{\text{gs}}$ target (closed neutron 1p-shell)
 direct 3n-transfer to the open sd-shell

\Rightarrow population of $(sd)^3$ structures expected ($3p-0h$)



Comparison of ^{17}C - ^{16}C spectra

Spectra aligned for the states with $(1d5/2)^3, 9/2^+$ configurations, in ^{16}C at 8.92 MeV and in ^{17}C at 3.10 MeV. (offset 5.82 MeV, same scale factor)

Red lines are almost parallel !
 connect states of corresponding structure
 \Rightarrow pairs of corresponding states

Comparison of observed excited states of ^{17}C and ^{16}C

State No.	2	3	4	5	6	7	8	9	10	11
$E_x(^{17}\text{C})$ [MeV]	0.31	2.06	3.10	4.25	6.20	6.75	7.47	8.94	10.56	11.71

Comparison of observed excited states of ^{17}C and ^{16}C

State No.	2	3	4	5	6	7	8	9	10	11
$E_x(^{17}\text{C})$ [MeV]	0.31	2.06	3.10	4.25	6.20	6.75	7.47	8.94	10.56	11.71

$E_x(^{16}\text{C})$ [MeV]	6.11 ^a	7.74	8.92	9.98	11.85	12.54	13.12	14.90	16.44	17.4
$E_x(^{16}\text{C})$ - 5.82	0.29	1.92	3.10	4.16	6.03	6.72	7.30	9.08	10.62	11.58
$\Delta E_x(^{16}\text{C}^*, ^{17}\text{C})$	-0.02	-0.14	± 0.0	-0.09	-0.17	-0.03	-0.17	+0.14	+0.06	-0.13

Comparison of observed excited states of ^{17}C and ^{16}C

reference from strongest states with stretched configurations $[(5/2^+)^3] 9/2^+$

State No.	2	3	4	5	6	7	8	9	10	11
$E_x(^{17}\text{C})$ [MeV]	0.31	2.06	3.10	4.25	6.20	6.75	7.47	8.94	10.56	11.71
$E_x(^{16}\text{C})$ [MeV]	6.11 ^a	7.74	8.92	9.98	11.85	12.54	13.12	14.90	16.44	17.4
$E_x(^{16}\text{C}) - 5.82$	0.29	1.92	3.10	4.16	6.03	6.72	7.30	9.08	10.62	11.58
$\Delta E_x(^{16}\text{C}^*, ^{17}\text{C})$	-0.02	-0.14	± 0.0	-0.09	-0.17	-0.03	-0.17	+0.14	+0.06	-0.13

Comparison of observed excited states of ^{17}C and ^{16}C

reference from strongest states with stretched configurations $[(5/2^+)^3] 9/2^+$


State No.	2	3	4	5	6	7	8	9	10	11
$E_x(^{17}\text{C})$ [MeV]	0.31	2.06	3.10	4.25	6.20	6.75	7.47	8.94	10.56	11.71
$E_x(^{16}\text{C})$ [MeV]	6.11 ^a	7.74	8.92	9.98	11.85	12.54	13.12	14.90	16.44	17.4
$E_x(^{16}\text{C}) - 5.82$	0.29	1.92	3.10	4.16	6.03	6.72	7.30	9.08	10.62	11.58
$\Delta E_x(^{16}\text{C}^*, ^{17}\text{C})$	-0.02	-0.14	± 0.0	-0.09	-0.17	-0.03	-0.17	+0.14	+0.06	-0.13
$\Gamma(^{17}\text{C})$ [MeV]	-	0.25	0.10	0.14	0.35	(0.20)	0.50	0.60	0.30	0.30
$\Gamma(^{16}\text{C})$ [MeV]	<0.03 ^a	0.20	≤ 0.10	0.12	0.22	(0.20)	0.40	0.30	0.15	0.15
$d\sigma(^{16}\text{C})/d\sigma(^{17}\text{C})$	-	2.9	2.0	2.9	3.3	3.0	1.7	0.6	1.3	1.6

Comparison of observed excited states of ^{17}C and ^{16}C

reference from strongest states with stretched configurations $[(5/2^+)^3] 9/2^+$

State No.	2	3	4	5	6	7	8	9	10	11
$E_x(^{17}\text{C})$ [MeV]	0.31	2.06	3.10	4.25	6.20	6.75	7.47	8.94	10.56	11.71
$E_x(^{16}\text{C})$ [MeV]	6.11 ^a	7.74	8.92	9.98	11.85	12.54	13.12	14.90	16.44	17.4
$E_x(^{16}\text{C}) - 5.82$	0.29	1.92	3.10	4.16	6.03	6.72	7.30	9.08	10.62	11.58
$\Delta E_x(^{16}\text{C}^*, ^{17}\text{C})$	-0.02	-0.14	± 0.0	-0.09	-0.17	-0.03	-0.17	+0.14	+0.06	-0.13
$\Gamma(^{17}\text{C})$ [MeV]	-	0.25	0.10	0.14	0.35	(0.20)	0.50	0.60	0.30	0.30
$\Gamma(^{16}\text{C})$ [MeV]	<0.03 ^a	0.20	≤ 0.10	0.12	0.22	(0.20)	0.40	0.30	0.15	0.15
$d\sigma(^{16}\text{C})/d\sigma(^{17}\text{C})$	-	2.9	2.0	2.9	3.3	3.0	1.7	0.6	1.3	1.6

Conclusion: These pairs of corresponding states have a common $3n$ -structure. Direct population of $(sd)^3$ structures on ^{14}C , ^{13}C through the reaction mechanism. Change of the core changes the excitation energies only by a *global constant*.



Code OXBASH
WBP interaction

^{17}C model space:

even-parity states


neutrons:

3 active neutrons in the (sd)-shell


4 active protons in the (psd)-shell,

proton 2⁺ core excitation was allowed, but the excitation needs 2nd order process and more than 7 MeV extra excitation energy: small cross sections expected (these are not displayed here, all other states up to 9.5 MeV are shown).


^{17}C (sd)³ three-neutron occupancies: OXBASH calculations





^{17}C (sd)³ three-neutron occupancies: OXBASH calculations



configurations	$3/2^+$	$1/2^+$	$5/2^+$	$7/2^+$	$9/2^+$	$11/2^+$
(1d5/2) ³	32%		47%		39%	
(1d5/2) ² (2s1/2) ¹	31%	65%		65%	16%	
(1d5/2) ² (1d3/2) ¹	2%		3%	3%	8%	68%
(1d5/2) ¹ (2s1/2) ²			19%			
(1d5/2) ¹ (2s1/2) ¹ (1d3/2) ¹	6%	2%		7%	10%	
(1d5/2) ¹ (1d3/2) ²			5%			
(2s1/2) ¹ (1d3/2) ²		5%				
small components	9%	5%	6%	7%	3%	8%
$\pi(1p3/2) 3(1p1/2) 1$	10%	12%	9%	10%	13%	14%
$\pi(1p3/2) 2(1p1/2) 2$	10%	11%	10%	8%	11%	10%



Decay properties, one-neutron decay



known (2p - xh) structures, (x= 0, 1, 2) from (t,p) in ^{16}C , ^{15}C , ^{14}C

Reduced decay energies :
 $E_{\text{dec}} = E_{x,i} - S_n - E_{x,f}$
 due to n-decay from (3p - xh) to (2p - xh) structures
 $\Rightarrow \Gamma_{\text{exp}}(E_{\text{dec}})$

Spectroscopic information: reduced widths $\gamma_{\ell, \text{exp}}^2$

n-decay



$$\Gamma_{\text{exp}} = 2 \gamma_{\ell, \text{exp}}^2 P_{\ell}(E_{\text{dec}})$$

^{17}C	E_x ^{17}C	E_x (MeV) $\rightarrow ^{16}\text{C}, J^{\pi}$	E_{dec} (MeV)	Γ_{exp} (MeV)	$\gamma_{\ell, \text{exp}}^2$ (MeV)	
					$\ell=0$	$\ell=2$
(3/2 ⁺)	2.06	$\rightarrow 0.00, 0^+$	1.33	0.25	0.085	0.37
9/2 ⁺	3.10	$\rightarrow 1.77, 2^+$	0.60	0.08	0.041	0.55
(5/2 ⁺ , 9/2 ⁺)	4.25	$\rightarrow 1.77, 2^+$	1.75	0.14	0.042	0.13
(5/2 ⁺)	6.20	$\rightarrow 3.03, 0^+$	2.44	0.35	0.088	0.21
(5/2 ⁺ , 7/2 ⁺)	6.75	$\rightarrow 3.98, 2^+$	1.49	(0.2)	0.110	0.43
(11/2 ⁺)	7.47	$\rightarrow 3.98, 2^+$	2.04	(0.2)	0.055	0.15
(11/2 ⁺)	7.47	$\rightarrow 4.14, 4^+$	2.60	0.50	0.120	0.27
^{16}C						
^{16}C	^{16}C	$\rightarrow ^{15}\text{C}, J^{\pi}$				
(1 ⁻ - 3 ⁻)	7.74	$\rightarrow 3.10, 1/2^-$	0.39	0.15	0.094	2.69
5 ⁻	8.92	$\rightarrow 4.22, 5/2^-$	0.45	0.04	0.023	0.52
(3 ⁻ , 5 ⁻)	9.98	$\rightarrow 4.22, 5/2^-$	1.51	0.12	0.038	0.14
(3 ⁻)	11.85	$\rightarrow 5.87, 1/2^-$	1.73	0.22	0.066	0.21
(3 ⁻)	12.54	$\rightarrow 5.87, 1/2^-$	2.42	(0.2)	0.051	0.12
(6 ⁻ , 5 ⁻)	13.12	$\rightarrow 6.84, 9/2^-$	2.03	0.40	0.110	0.30

Summary and Conclusions

The structure and assignments of ^{17}C and ^{16}C were discussed using the

- **dependence of cross sections on dynamical matching conditions**
- **striking similarity observed in the level schemes of ^{16}C and ^{17}C , indicating a common structure of $(sd)^3$ -type for corresp. pairs of states**
- **changing the core, excitation energies change only by a global constant**
- **shell model calculations for ^{17}C , comparison to tentative assignments**
- **comparison to ^{19}O experimental levels and SM calculations**
- **exper. widths of resonances, decay properties, reduced widths**

The neutron $(sd)^3$ excitation energies observed in ^{17}C and ^{16}C seem to be almost completely independent (within ± 0.16 MeV) from the corresponding cores ^{14}C , ^{13}C , respectively, except a global constant. This is found for 10 pairs of states for ^{17}C , ^{16}C over a range of 10 MeV excitation energy.