Order α^2 corrections to the decay rate of orthopositronium

G.S. Adkins, R. Fell¹, and J. Sapirstein²

Department of Physics and Astronomy, Franklin & Marshall College
P.O. Box 3003, Lancaster, PA, 17604 USA
Tel +717-291-4245, Fax +717-399-4474
E-mail: g_adkins@acad.fandm.edu

1 Brandeis University, Waltham, MA, 01742 USA

² Department of Physics, University of Notre Dame, Notre Dame, IN, 46556 USA

The discrepancy between theory and experiment for the decay rate of orthopositronium has long been one of the outstanding problems in precision QED. The calculated decay rate is [1, 2, 3]

$$\Gamma_{\text{o-Ps}} = \Gamma_0 \left[1 + A \frac{\alpha}{\pi} + \frac{1}{3} \alpha^2 \ln \alpha + B \left(\frac{\alpha}{\pi} \right)^2 - \frac{3\alpha^3}{2\pi} \ln^2 \alpha + \cdots \right] ,$$

where the lowest order contribution is $\Gamma_0 = \frac{2}{9}(\pi^2 - 9)\frac{m\alpha^6}{\pi} = 7.211169\mu s^{-1}$ [4]. The one-loop correction is known to be A = -10.286606(10) [5]. The result of the present calculation is a value for the two-loop correction B.

Our calculation was done in the context of Nonrelativistic Quantum Electrodynamics (also known as NRQED) [6] following the approach outlined by Labelle, Lepage, and Magnea [7]. This method allows the high-energy part of the calculation to be treated as an on-shell scattering process. The high-energy calculation is part of a "matching" procedure in which a set of nonrelativistic interaction operators is defined. These operators are used to work out the bound-state aspects of the problem. Our calculation of the high-energy process followed by a bound-state calculation using the effective interaction operators allowed us to complete the determination of B [8].

Acknowledgments. The work of GA was partially supported by NSF grants PHY-9711991 and PHY-9722074, and that of JS by PHY-9870017. Useful conversations with P. Labelle, G.P. Lepage, and R. Hill are acknowledged.

- [1] W.E. Caswell, G.P. Lepage, and J. Sapirstein, Phys. Rev. Lett. 38, 488 (1977).
- [2] W.E. Caswell and G.P. Lepage, Phys. Rev. A 20, 36 (1979).
- [3] S.G. Karshenboim, Zh. Eksp. Teor. Fiz. 103, 1105 (1993) [JETP 76, 541 (1993)].
- [4] A. Ore and J.L. Powell, *Phys. Rev.* **75**, 1696 (1949).
- [5] G.S. Adkins, Phys. Rev. Lett. 76, 4903 (1996).
- [6] W.E. Caswell and G.P. Lepage, Phys. Lett. 167B, 437 (1986).
- [7] P. Labelle, G.P. Lepage, and U. Magnea, Phys. Rev. Lett. 72, 2006 (1994).
- [8] G.S. Adkins, R. Fell, and J. Sapirstein, hep-ph/0003028.