Relativistic and QED corrections to the metastable states of the antiprotonic helium atoms

Yasushi Kino, Nobuhiro Yamanaka¹, Masayasu Kamimura² and Hiroshi Kudo

Department of Chemistry, Tohoku University, Sendai 980-8578, Japan Tel +81-22-217-6596, Fax +81-22-217-6597

E-mail: kino@mail.cc.tohoku.ac.jp

Antiprotonic helium atom ($\bar{p}\mathrm{He^+}$) is an exotic three-body system consisting of an antiproton (\bar{p}), a helium nucleus ($\mathrm{He^{2+}}$) and an electron (e^-). The transition wavelengths were measured with precise laser spectroscopy (see [1], further references therein) for the metastable states. The data are very challenging to the few-body theory since it is necessary to accomplish very high precision in an energy-level calculation managing large interacting angular momenta of 30–40. Moreover the relativistic and QED effects exceed this accuracy [2]. To calculate the relativistic effects, we start with the Breit equation. Using Pauli approximation, we obtain the three-body Hamiltonian H for the Schödinger equation,

$$H = \sum_{i} \frac{\mathbf{p}_{i}^{2}}{2m_{i}} + \sum_{i>j} \frac{z_{i}z_{j}}{r_{ij}} - K_{cm} + \alpha^{2}H_{rel} + \alpha^{3}H_{QED}, \tag{1}$$

with

$$H_{\text{rel}} = -\sum_{i} \frac{\mathbf{p}_{i}^{4}}{8m_{i}^{3}} + \sum_{i>j} \left[-\frac{z_{i}z_{j}}{2} \left(\frac{1}{m_{i}^{2}} + \frac{1}{m_{j}^{2}} \right) \pi \delta(\mathbf{r}_{ij}) - \frac{z_{i}z_{j}}{2m_{i}m_{j}} \left(\frac{\mathbf{p}_{i} \cdot \mathbf{p}_{i}}{r_{ij}} + \frac{(\mathbf{r}_{ij} \cdot \mathbf{p}_{i})(\mathbf{r}_{ij} \cdot \mathbf{p}_{j})}{r_{ij}} \right) \right.$$

$$\left. + \frac{(z_{i} - 2\mu_{i})z_{j}}{2m_{i}^{2}} \frac{(\mathbf{r}_{ij} \times \mathbf{p}_{i}) \cdot \mathbf{s}_{i}}{r_{ij}^{3}} - \frac{z_{i}z_{j}}{m_{i}m_{j}} \frac{(\mathbf{r}_{ij} \times \mathbf{p}_{i}) \cdot \mathbf{s}_{j}}{r_{ij}^{3}} \right.$$

$$\left. - \frac{8\pi}{3} \frac{\mu_{i}\mu_{j}}{m_{i}m_{j}} (\mathbf{s}_{i} \cdot \mathbf{s}_{j}) \delta(\mathbf{r}_{ij}) + \frac{\mu_{i}\mu_{j}}{m_{i}m_{j}} \left((\mathbf{s}_{i} \cdot \mathbf{s}_{j}) - 3 \frac{(\mathbf{r}_{ij} \times \mathbf{s}_{i})(\mathbf{r}_{ij} \times \mathbf{s}_{j})}{r_{ij}^{5}} \right) \right], \quad (2)$$

$$H_{\text{QED}} = -\frac{4}{3} \left(\frac{19}{30} - 2 \ln \alpha + \ln K_{0} \right) \left(z_{\alpha} z_{e} \delta(\mathbf{r}_{\alpha e}) + z_{\bar{p}} z_{e} \delta(\mathbf{r}_{\bar{p}e}) \right), \quad (3)$$

where \mathbf{p}_i , m_i z_i , μ_i and \mathbf{r}_{ij} are the momentum of *i*th particle, its mass, its electric charge, its magnetic dipole moment and the relative vector between *i*th particle and *j*th particle. The value $\ln K_0 = 4.4$ is the Bethe logarithm. We subtract the kinetic operator $K_{\rm cm}$ of the center of mass system from the Hamiltonian. The operator $H_{\rm QED}$ corresponding to the one-loop vacuum polarization and self-energy is added to the Hamiltonian. The total wave function is given by

$$\Phi_{vJFM} = \sum_{K} C_{K} \left[\left[\Psi_{vJ}^{\bar{p}\text{He}^{\dagger}} \otimes \xi_{e} \right]_{K} \otimes \xi_{\bar{p}} \right]_{FM}, \tag{4}$$

¹ Department of Physics, University of Tokyo, Tokyo 113-0033, Japan

² Department of Physics, Kyushu University, Fukuoka 812-8581, Japan

where $\Psi_{vJ}^{\bar{p}\mathrm{He}^+}$ is a non-relativistic three-body wave function, C_K is an expansion coefficient and ξ is a spin function. The non-relativistic wave functions and energies are calculated with a non-adiabatic coupled rearrangement channel method [3] diagonalizing the non-relativistic part of the Hamiltonian. The three rearrangement channels are shown in Ref. [4]. The relativistic and QED corrections are calculated with the first order perturbation theory. In Table 1, the transition wavelengths and uncertainty of the antiproton mass (charge) are listed. The relativistic and QED collection reduced the discrepancy between the calculated and observed values by about 40 ppm. To estimate the uncertainty of \bar{p} mass from the experimental data, we recalculated the wavelength using the \bar{p} mass and charge scaled with $1 + \Delta x$ against the p mass and charge, $m_{\bar{p}} = (1 + \Delta x)m_p$ and $e_{\bar{p}} = (1 + \Delta x)e_p$. The value Δx was estimated using an uncertainty of the experimental wavelengths $\Delta \lambda_{\rm expt}$,

$$\Delta x = \frac{\Delta m_{\bar{p}}}{m_p} = \Delta \lambda_{\text{expt}} \left(\frac{d\lambda}{dx}\right)_{\text{col}}^{-1}.$$
 (5)

We reduced uncertainty of the \bar{p} mass (or equivalently charge) by two orders of magnitude [5]. We also calculated the fine and hyperfine splitting of the metastable states [7] for the forthcoming experiment using microwave resonance at CERN. The values agree well with other calculation [6]. We obtained the transition wavelengths 713.5819 nm for $(J=35, v=1, K=J+1/2) \rightarrow (34, 3, J+1/2)$ and 713.5812 nm for $(J=35, v=1, K=J-1/2) \rightarrow (34, 3, J-1/2)$, and obtained the transition frequencies 12.899 GHz for $(J=35, v=1, K=J-1/2, F=J) \rightarrow (35, 1, J+1/2, J+1)$ and 12.927 GHz for $(J=35, v=1, K=J-1/2, F=J-1) \rightarrow (35, 1, J+1/2, J)$.

Table 1: Transition wavelengths, λ_{NR} : pure Coulomb calculation, λ_{C} : calculation with relativistic and QED corrections and λ_{E} : experimental values. Δx is the uncertainty of \bar{p} mass.

$\overline{(J_i,v_i) – (J_f,v_f)}$	$\lambda_{ m NR} \ (m nm)$	$\lambda_{ m C} \ ({ m nm})$	$\lambda_{ m E} \left[1 ight] \left({ m nm} ight)$	Δx
$\overline{(35,3)}$ – $(34,3)$	597.2290	597.2573	597.2570 ± 0.0003	1×10^{-7}
(34,2)- $(33,2)$	470.7048	470.7220	470.7220 ± 0.0006	3×10^{-7}

Acknowledgments. This work is financially supported by the Grant-in-Aid for Scientific Research, Ministry of Education and Culture. YK wish to acknowledge Inoue Foundation for Science for their generous financial support. The computations were partially carried out on the RIKEN supercomputer VPP700/128.

- [1] H.A. Torii et al., Phys. Rev. A59, 223 (1999).
- [2] V.I. Korobov and D.D. Bakalov, Phys. Rev. Lett. 79, 3379 (1997).
- [3] M. Kamimura, Phys. Rev. A 38, 621 (1988).
- [4] Y. Kino M. Kamimura and H. Kudo, Nucl. Phys. A 631, 649 (1998).
- [5] Y. Kino M. Kamimura and H. Kudo, Hyp. Interact. 119, 201 (1999).
- [6] D. Bakalov and V.I. Korobov, Phys. Rev. A 57, 1662 (1998).
- [7] N. Yamanaka, Y. Kino M. Kamimura and H. Kudo, to be published.