Dark-hollow beam guiding of a continuous low-velocity atomic beam

Min Yan, Jianping Yin¹, Yifu Zhu

Department of Physics, Florida International University
University Park, Miami, FL 33199, USA
Tel 305-348-4075, Fax 305-348-6700
E-mail: yifuzhu@fiu.edu, Website: http://www.fiu.edu/ yifuzhu

1 Department of Physics, Suzhou University, Suzhou, China

Optical beams with zero central intensity can be called dark hollow beams (DHB) and have been used in various studies of neutral atom manipulation in recent years. In a blue detuned DHB, atoms experience a dipole force towards the beam center and can be confined along the beam under appropriate conditions. Several recent experiments have demonstrated atom guiding and atom transverse cooling in a DHB [1, 4]. Possibilities of atom guiding and Sisyphus cooling in a DHB have been discussed in the literature [5].

We report experimental observations of dark-hollow-beam guiding, focusing, and splitting of a continuous, low velocity atomic beam generated in a magneto optical trap. Our experiments were carried out in a standard 87Rb magneto-optical trap (MOT) and a low-velocity rubidium atomic beam was generated by placing a 1 mm black spot in the center of the vertical returning trapping and cooling beam as first demonstrated by Lu et al [6]. Two extended-cavity diode lasers were used as the trapping laser and repump laser respectively. With a simple optical axicon set up [7], a Ti:Sapphire laser beam was converted into a DHB of power 250 mW.

The loading and guiding characteristics of the low-velocity Rb beam can be manipulated by beam shaping of the DHB. We varied the dark center diameter of the DHB in a range of 0.1 mm to 1 mm and moved the DHB waist above or below the MOT. Our results show that when the DHB waist is below the MOT, the guided atomic beam converges; when the DHB waist is above the MOT, the guided atomic beam diverges.

Acknowledgment. This work is supported by ONR and ARO.

- [1] T. Kuga et al, Phys. Phys. Lett. 78 4713 (1997).
- [2] Y. Ovchinnikov et al, Phys. Phys. Lett. 79 2225 (1997).
- [3] S. Kuppens et al, Phys. Phys. A 58 3068 (1998).
- [4] Xu et al, Phys. Phys. A 60 4796 (1999).
- [5] J. Yin et al, J. Opt. Soc. Am. B 15 47132235 (1998).
- [6] Z. T. Lu et al, Phys. Phys. Lett. 77 3331 (1996).
- [7] I. Manek et al, Opt. Commun. 147 67 (1998).